

Crossover from Spin-Density-Wave to Neel-like Ground State

Jaime Ferrer

Departamento de Física de la Materia Condensada, Universidad Autónoma de Madrid, 28049

Madrid, Spain

Abstract

The characterization and evolution of a Spin Density Wave into the Quantum Neel ground state is considered in the context of a weak coupling theory of the half-filled Hubbard model. Magnetic properties obtained from this weak coupling approach in one dimension compare favorably with exact results from Bethe ansatz (BA). A study of the evolution of several length scales from weak to strong coupling is also presented.

71.27.+a, 75.10.Lp, 75.30.Fv

Typeset using REVTeX

I. INTRODUCTION

Interest in low-dimensional Quantum Antiferromagnetism experienced a sharp rise with the realization that magnetic fluctuations are at the root of many of the exotic properties of High Temperature Superconductors. It is fairly well established by now that the plain two-dimensional Quantum Heisenberg model can account for most of the experimental features of undoped cuprates¹⁻³. It is also widely believed that the simple one-band Hubbard Model (HM) can map large parts of the experimental phase diagram^{1,4-6}, and most of the theoretical work has used it as an appropriate starting point^{2,3}. Simple as it looks, the HM displays a rich variety of regimes, with highly non-trivial physics. In fact, there is no consensus about the adequate low-energy effective action for small doping concentrations⁷⁻⁹, nor about the gross features of the phase diagram in that regime^{7,10,11}.

The situation is much more clear exactly at half-filling, because charge degrees of freedom are frozen out. In this case and for large Coulomb repulsion, the physical electrons are localized, and the ground state is the quantum analog of the Neel state. In the opposite limit of small U , the ground state is a Spin Density Wave (SDW), and should be adequately described using standard RPA theory over a broken symmetry, Mean Field SDW state. In an early development, Schrieffer, Wen and Zhang¹² proposed that the later, weak coupling, approach (to be called RPA-SDW hereafter) could also explain the physics of the Quantum Neel state. Later on, Chubukov and Frenkel¹³ explicitly showed that many of the known results for the Heisenberg model could indeed be recovered within the SDW-RPA approach. There are, nevertheless, some details missing in the physical picture of the half-filled HM. For instance: it is common lore that a SDW is the analog of the Neel state when the electrons are itinerant, but a clear and quantitative description of it is lacking. As a direct consequence, the crossover regime between both states is not fully understood. The author also feels that there is some confusion about the Temperature and length scales generated in the different regimes. To be specific: these scales are, in general, well established for the positive- U HM at strong coupling and for the negative- U case at weak coupling, but not in the other two

limits¹⁴.

The purpose of this paper is to shed some more light on these issues. In order to do so, I will apply the RPA-SDW expansion used in ref. 12 to the one-dimensional case at Zero Temperature and compare with the Bethe ansatz solution¹⁵. Hence, the whole phase diagram at half-filling will be covered. I will also draw many consequences on the scales involved in the problem by the heavy use of duality relations¹⁶. The computations are performed for the worst case, because RPA schemes are supposed to do a bad job in low dimensions and in the spin-disordered phase. But I will actually show that RPA works remarkably well even in 1d -at half filling-.

Some comments are in order at this point of the introduction. The HM at half-filling falls into the $O(3)$, instead of XY, universality class. A closely related fact is that its ground state is a singlet in the strong coupling limit¹⁷ so that all correlation functions are rotationally invariant. A further, specific feature of one-dimensional systems is that they do not possess Long Range Order, even at zero Temperature, because of thermal and quantum fluctuations¹⁸. Correlation functions related with the order parameter have a mass gap and do not follow Goldstone behavior¹⁹. The complications derived from spin-charge separation are eluded because charge degrees of freedom are frozen out at half-filling¹⁵.

The layout of this paper is as follows: several magnitudes will be computed within RPA-SDW and compared with Bethe ansatz results in section 2. Section 3 will present a study of the different length and Temperature scales of the problem. A brief conclusion will end the paper. Details of the calculations will be relegated to appendices A and B. Energies will be measured in units of the tunneling amplitude, t .

II. STUDY OF THE SDW-RPA ANSATZ AT $T = 0$

It is commonly believed that RPA approaches should not perform well in low dimensions or when one is trying to describe physical properties of the strong coupling phase. The purpose of this section is to show that RPA-SDW produces accurate results at either weak

or strong coupling even in the extreme case of one dimension, as long as the ground state is close enough to a commensurate Antiferromagnet.

The first quantity which arises in RPA-SDW theory is the magnitude of the Mean Field charge gap, 2Δ , which is obtained from the equation

$$\frac{1}{U} = \frac{1}{N} \sum'_k \frac{1}{E_k} \quad (1)$$

Inclusion of RPA fluctuations does not modify the equation. Although the asymptotic expression for Δ when U approaches zero, $\frac{8}{\pi} U e^{-2\pi/U}$, is not captured by the RPA-SDW approximation, which gives $\Delta \sim e^{-2\pi/\sqrt{U}}$, the numerical differences are negligible for such small values of U (see fig. 1). The RPA-SDW curve begins to depart appreciably from the exact result for values of U of the order of the bandwidth, D . This fact suggests that the crossover from nearly itinerant to localized electron behavior is not described quantitatively within this weak-coupling approximation.

The exact large- U value of the charge gap, on the other hand, is $U - 4t = U - D$, where D is the free band width²⁰. RPA-SDW theory overestimates 2Δ by D , because the approximate lower and upper Hubbard bands tend to δ -functions while the exact ones retain a bandwidth of D when $U \rightarrow \infty$ (see fig. 1 again).

The collective excitations of the system appear as poles of the imaginary part of the different correlation functions $\chi''(q, w)$. The weight of a particular fluctuation (q, w) on the ground state is given by the residue of χ'' at that particular value of (q, w) . The RPA-SDW ansatz, in conjunction with Goldstone's theorem, tells us that the only low-energy collective excitations of the half-filled HM at zero Temperature and for large U are spin waves, and show up as poles of the transverse spin susceptibility, χ_{RPA}^{+-} . Charge and longitudinal spin fluctuations correspond to poles of χ_{RPA}^{00} and χ_{RPA}^{zz} , and have a mass gap. These correlation functions have a very simple form in the strong coupling limit. For instance

$$\chi_{RPA}^{+-}(q, w) = \frac{1}{U} \frac{w^2 - w_q^2 + 4(1 - \cos(q))}{2w_q} \left\{ \frac{1}{w + w_q + i\delta} - \frac{1}{w - w_q + i\delta} \right\} \quad (2)$$

Fig. 2 (a) shows the imaginary part of the transverse spin susceptibility for $U = 20$. It displays the typical Goldstone-mode structure and has a strong q -dependence, proper for

spin waves. Notice also that the weight of the Ferromagnetic zero mode tends to zero, while that of the Antiferromagnetic one diverges, in agreement with strong coupling analyses²¹. On the contrary, $\chi_{RPA}^{zz}(q, w)$ for $U = 20$ is almost dispersionless and has a large gap of order U (see figure 2 (b)). Hence, longitudinal spin fluctuations are massive and possess a localized nature.

The exact spin correlation functions should be rotationally invariant because the ground state of the system is a singlet at strong coupling. Regretfully, the RPA-SDW response functions do not have this property.

Returning to the strong-coupling case again, it is easy to show that the excitation spectrum given by RPA-SDW is

$$w_q = v_s |\sin(q)| \sim_{q \rightarrow 0} v_s q \quad (3)$$

This is the result one would obtain for the Heisenberg model in Linear Spin Wave theory (LSWT)¹⁷. The linear relation between w and q for small q holds all the way down to $U = 0$ and serves to define v_s for all values of the Coulomb repulsion (fig. 3). Notice that the RPA-SDW velocity grows as U increases from zero and then has a maximum for $U \sim 1$. What happens here is that the limit $U \rightarrow 0$, where $v_s = 2$, does not coincide with the $U = 0$ case ($v_s = 0$). This subtle point is well captured by the exact BA analysis, but not by RPA-SDW theory, where v_s begins at zero, increases steeply until it reaches a value close to 3 at the maximum and then approaches the BA curve for $U \sim D$. Both curves decrease approximately at the same pace for even larger U and tend asymptotically to $\frac{\pi}{2}J \sim 1.6J$ and J . The value given by RPA-SDW theory for v_s in higher dimensions ($\sqrt{2}J$ for $d=2$) coincides with that obtained from LSWT.

The squared magnetic moment, $\langle S^2 \rangle$ is displayed in fig. 4, as given by Mean Field Theory (dashed line) and Bethe ansatz (solid line). It is quite remarkable the fact that Mean Field Theory already give the right $U \rightarrow 0$ and $U \rightarrow \infty$ limits. Notice also how Mean Field Theory underestimates grossly $\langle S^2 \rangle$ for small values of U , while tends correctly to $3/4$ in the opposite limit. What this seems to imply again is that the Mean Field spin degrees

of freedom are *too itinerant* in the crossover regime from itinerant to localized electron behavior.

RPA fluctuations do not cure this flaw. In fact, $\langle S^2 \rangle$ diverges at strong coupling as

$$\langle S^2 \rangle = \frac{1}{4} \sum_k \frac{1 - \cos(k)}{\sin(k)} \rightarrow \infty \quad (4)$$

which is also the result obtained by LSWT. In the opposite limit, on the contrary, RPA-SDW theory reproduces the exact free result $\frac{3}{8}$.

Another important result concerns the average of the staggered magnetization for large U , which also coincides with LSWT¹³

$$\langle S^z \rangle = \frac{1}{2} \left\{ 1 - 2 \sum_k \frac{1 - \sin(k)}{|\sin(k)|} \right\} \rightarrow -\infty \quad (5)$$

This result implies that one can obtain a charge insulating spin disordered ground state, also called *spin liquid*, within SDW-RPA theory, because the charge gap is still given by the Mean Field result. LSWT is known to diverge in one dimension due to infrared divergences, but to lead to amazingly accurate results in higher dimensions^{22,23}.

Figure 5 is a useful eye-guide to understand the differences between the Neel state and a Spin Density Wave. For large values of U , the charge degrees of freedom of physical electrons are frozen out and the spin degrees of freedom become localized magnetic moments: the Heisenberg Hamiltonian provides a good description of the HM in this limit. In this case, $\langle S^2 \rangle = S(S + 1) = 3/4$. In the opposite case, $\langle S^2 \rangle$ equals $3/8$, which is the value for free electrons. As soon as U differs from zero, a charge gap opens up, and the expectation value of the staggered magnetization –in dimensions higher than one– becomes finite albeit small. The squared magnetic moment increases with U from $3/8$ to $3/4$ continuously but with a steep slope. By the time when U equals the band width, $\langle S^2 \rangle$ is already very close to $3/4$, and the Spin Density Wave has fully developed into the Neel state. Likewise, for that value of U , the exact spin wave velocity almost coincides with $\frac{\pi}{2}J$.

III. LENGTH SCALES IN THE HUBBARD MODEL

The transformation

$$\begin{aligned}\hat{f}_{i\uparrow}^+ &\Leftrightarrow \hat{f}_{i\uparrow}^+ \\ \hat{f}_{i\downarrow}^+ &\Leftrightarrow e^{i\mathbf{Q}\mathbf{R}_i} \hat{f}_{i\uparrow}^+\end{aligned}\tag{6}$$

where $\mathbf{Q} = (\pi, \pi, \dots)$, changes the positive- U half filled HM into the negative one, and vice versa. It also exchanges spin and isospin operators¹⁶:

$$\begin{aligned}\hat{\mathcal{H}} = -t \sum_{\langle i,j \rangle, \sigma} \hat{f}_{i,\sigma}^+ \hat{f}_{j,\sigma} + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} &\Leftrightarrow \hat{\mathcal{H}} = -t \sum_{\langle i,j \rangle, \sigma} \hat{f}_{i,\sigma}^+ \hat{f}_{j,\sigma} - U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} \\ \left\{ \frac{1}{2}(\hat{n}_{i\uparrow} - \hat{n}_{i\downarrow}), e^{-i\mathbf{Q}\mathbf{R}_i} \hat{f}_{i\uparrow}^+ \hat{f}_{i\downarrow}, e^{i\mathbf{Q}\mathbf{R}_i} \hat{f}_{i\downarrow}^+ \hat{f}_{i\uparrow} \right\} &\Leftrightarrow \left\{ \frac{1}{2}(\hat{n}_i - 1), \hat{f}_{i\uparrow}^+ \hat{f}_{i\downarrow}^+, \hat{f}_{i\downarrow} \hat{f}_{i\uparrow} \right\}\end{aligned}$$

$$(\hat{S}^z, e^{-i\mathbf{Q}\mathbf{R}_i} \hat{S}^+, e^{i\mathbf{Q}\mathbf{R}_i} \hat{S}^-) \Leftrightarrow (\hat{\psi}^z, \hat{\psi}^+, \hat{\psi})\tag{7}$$

These duality relations allow to infer many properties of one model from the other. For instance, the positive- U HM at half-filling possesses a global $SU(2)_{sp} \otimes U(1)_{ch}$. This means that its counterpart has a global $SU(2)_{ch} \otimes U(1)_{sp}$. Likewise, an

- Antiferromagnet (AFM) polarized along the Z-axis is equivalent to a commensurate Charge Density Wave
- AFM polarized along the XY-plane is equivalent to a uniform superconductor
- Ferromagnet (FM) polarized along the Z-axis is equivalent to the uniform state
- FM polarized along the XY-plane is equivalent to a commensurate *staggered* superconductor²⁴

Because the ground state of an AFM at strong coupling is a singlet, a uniform superconductor should be degenerate with a CDW for large U . Conversely, the spin transport properties of an AFM polarized along the Z-axis and XY-plane should be very different, because the charge transport properties of a uniform superconductor and a commensurate CDW indeed are. The appearance of the nesting wavevector $\mathbf{Q}$ in the duality relations introduces, however, a slight difference because staggering means folding of the Brillouin zone into two pieces, with the corresponding doubling of the periodicity. It is also interesting to notice that there do not seem to be many experimental realizations of a staggered superconductor, while a uniform FM polarized in the XY-plane takes place in Nature very frequently²⁵.

The remaining of this section is devoted to elucidate what are the different length scales which appear in the positive and negative-U HM and how they show up in the following correlation functions:

$$\begin{aligned} \tilde{G}_{SDW}(k, k+Q) &= \begin{pmatrix} \langle T \hat{f}_\sigma(k) \hat{f}_\sigma^+(k) \rangle & \langle T \hat{f}_\downarrow(k) \hat{f}_\uparrow^+(k+Q) \rangle \\ \langle T \hat{f}_\uparrow(k+Q) \hat{f}_\downarrow^+(k) \rangle & \langle T \hat{f}_\sigma(k+Q) \hat{f}_\sigma^+(k+Q) \rangle \end{pmatrix} \Leftrightarrow \\ \tilde{G}_{sup}(k) &= \begin{pmatrix} G_0 & F \\ F^+ & G_0 \end{pmatrix} = \begin{pmatrix} \langle T \hat{f}_\sigma(k) \hat{f}_\sigma^+(k) \rangle & \langle T \hat{\psi}(k) \rangle \\ \langle T \hat{\psi}^+(k) \rangle & \langle T \hat{f}_{-\sigma}^+(k) \hat{f}_{-\sigma}^+(k) \rangle \end{pmatrix} \\ C^{+-}(k) &= \langle T S^+(k) S^-(-k) \rangle \Leftrightarrow C_\psi(k) = \langle T \psi^+(k) \psi(-k) \rangle \end{aligned} \quad (8)$$

The language proper for superconductivity will be used, and it is assumed that the dimension is larger than 2. RPA-SDW makes the ansatz that the spin is polarized along the Z-axis. However, I have chosen in this section to write down Green-functions which correspond to polarization in the XY plane, Eq. 8, to make duality arguments more explicit.

Generically, there are three different Temperature scales present in this problem, as shown schematically in figure 5. The highest Temperature, denoted as T_b , is that at which binding of uncorrelated pairs occurs. It is a crossover Temperature, signaled by a divergence in the T-matrix^{26,27}. A spin gap, $m_1 = 2\Delta$, opens up in G_0 because it costs some energy

to break a pair. Conversely, for U positive, a charge gap appears in G_0 . The non-diagonal elements of $\tilde{G}_{sup}$ and $\tilde{G}_{SDW}$, $F(k) = \langle T\hat{\psi}^+(k) \rangle$, are still zero.

At a lower temperature, T_0 ($\sim J$ for strong coupling), correlations among Cooper pairs build up, and the correlation length $\xi(T)$ begins to be longer than the lattice spacing, a . Its inverse, $m_{corr} = 1/\xi$, appears as a pole in the correlation function C_ψ , which takes the conventional Ornstein-Zernike form

$$C_\psi(k) = \frac{1}{\pi} \frac{m_{corr}}{k^2 + m_{corr}^2} \quad (9)$$

m_{corr} is the gap for excitations of the collective modes of the order parameter. For U positive, the pole appears in the correlation function C^{+-} , and m_{corr} is the gap in the spectrum of spin waves²⁸. This is also the Temperature where the anomalous Nambu Green's functions, $F(k \sim 1/a) = \langle T\hat{\psi}^+(k \sim 1/a) \rangle$ begin to differ from zero.

Finally, at a still lower temperature, T_c , a phase transition towards the broken symmetry superconducting state occurs, and is signaled by a divergence of $\xi(T)$: the mass gap m_{corr} closes down, as dictated by Goldstone's theorem. Below T_c , another length scale sets in: the coherence length of Cooper pairs, which is also Josephson's coherence length, $\xi_J = 1/m_J$ ²⁹:

$$C_\psi(k) = 4\pi\xi_J \frac{|\psi^*|^2}{k^2} = \frac{2K_B T}{\rho_s} \frac{|\psi^*|^2}{k^2} \quad (10)$$

where $\psi^* = \langle \hat{\psi}^+(k=0) \rangle$ is the expectation value of the order parameter and ρ_s is the superfluid density.

Notice that for small U , T_b , T_0 and T_c merge into one line; m_1 is then called the superconducting charge gap, and equals m_J for $T < T_c$, or m_{corr} for $T > T_c$. This means that $\xi(T)$ decreases to zero very quickly as the Temperature is raised above T_c in the weak coupling regime. In other words, Cooper pairs become completely uncorrelated for Temperatures slightly above T_c . Mean Field BCS theory is very accurate in this limit and predicts that the order parameter $\psi = \frac{m_1}{2U}$.

For larger attraction U , these three Temperatures depart from each other, and there is a finite Temperature range above T_c where $\xi(T)$ is different from zero. The order parameter

is no longer equal to $\frac{m_1}{2U}$ because fluctuations become important (remember the discussion in the former section).

Finally, for very large U , T_b diverges, and Cooper pairs become composite bosons. T_0 and T_c again go to zero because it costs more and more energy for a pair to hop from one site to the following -the effective tunneling amplitude, t^2/U , vanishes-. For the positive- U case, we would say that the spins become progressively less correlated because the exchange integral, $J \sim 4t^2/U$ vanishes.

IV. CONCLUSIONS

This paper shows that the RPA-SDW expansion developed in ref. 12 provides reliable results for the magnetic properties of the Hubbard model at either weak or strong coupling even in the extreme case of one spatial dimension. RPA-SDW does provide a qualitative, yet not quantitative, description of the intermediate coupling regime, when the Coulomb repulsion U is of the order of the bandwidth D . These conclusions should be quite robust against doping and Temperature as long as the exact ground state is still Neel-like^{7,10,11}. The accuracy of this expansion should improve with increasing dimensionality. The Spin Density Wave and Quantum Neel ground states are characterized the expectation value $\langle S^2 \rangle$, which measures the degree of itinerancy of the charge degree of freedom electrons. This criterion allows to show that the Spin Density Wave evolves continually into the Neel state. Free electrons have $\langle S^2 \rangle = 3/8$, but $\langle S^2 \rangle$ increases quickly with U and saturates to the value of localized spins, $3/4$, when the Coulomb energy U is of the order of the bandwidth. Several Temperature and length scales are discussed in both weak and strong coupling limits.

It is a pleasure to thank G. Gómez-Santos, N. Lorente, J. Pérez-Conde, F. Gebhard, F. Sols and E. Miranda for their helpful hints. This work was supported by DGICYT, Project No. PB93-1248

APPENDIX A: REMINDER OF SDW-RPA FORMALISM AT HALF-FILLING AND ZERO TEMPERATURE

The Hubbard Hamiltonian can be rewritten in terms of the staggered magnetization, $\hat{m}_i$

$$\hat{\mathcal{H}} = \sum_k \epsilon_{k,\sigma} \hat{f}_{k,\sigma}^+ \hat{f}_{k,\sigma} - U \sum_i \hat{m}_i^2 = \hat{\mathcal{H}} = -2t \sum_{k,\sigma} \cos(k) \hat{f}_{k,\sigma}^+ \hat{f}_{k,\sigma} - U \sum_i \hat{m}_i^2 \quad (\text{A1})$$

if we neglect charge fluctuations, a good assumption for the half-filled case. The staggered magnetization can, in turn, be expressed in terms of spin-1/2 operators

$$\begin{aligned} \hat{S}_i^\mu &= (\hat{\rho}_i, \hat{S}_i^x, \hat{S}_i^y, \hat{S}_i^z) = \sum_{\alpha,\beta} \hat{f}_{i,\alpha} \tau_{\alpha\beta}^\mu \hat{f}_{i,\beta} \\ \tau_{\alpha\beta}^\mu &= \left(\delta_{\alpha\beta}, \frac{\sigma_{\alpha\beta}^i}{2} \right) \end{aligned} \quad (\text{A2})$$

as $\hat{m}_i = e^{i\mathbf{Q}\cdot\mathbf{R}_i} \hat{S}_i^z$. A Mean Field decomposition on the interaction term is performed now, $\hat{m}_i^2 = \langle \hat{m}_i \rangle \hat{m}_i$, so that

$$\hat{\mathcal{H}} = \sum_{k,\sigma} \epsilon_k \hat{f}_{k,\sigma}^+ \hat{f}_{k,\sigma} + U \langle \hat{m}_i \rangle \sum_i \hat{m}_i \quad (\text{A3})$$

where $U \langle \hat{m}_i \rangle = \Delta$ is half the charge gap. The Hamiltonian is diagonalized with a conventional Bogoliubov transformation and the result is

$$\begin{aligned} \hat{\mathcal{H}} &= - \sum_k' E_k v_k^2 + \sum_{k,\sigma}' E_k (\hat{c}_{k,\sigma}^+ \hat{c}_{k,\sigma} - \hat{d}_{k,\sigma}^+ \hat{d}_{k,\sigma}) \\ E_k &= \sqrt{\epsilon_k^2 + \Delta^2} \end{aligned} \quad (\text{A4})$$

with the self-consistency condition

$$\frac{1}{U} = \frac{1}{N} \sum_k' \frac{1}{E_k} \quad (\text{A5})$$

The prime in the sum means summation in the restricted magnetic Brillouin zone. The new ground state, $|\mathcal{SDW}\rangle$, is annihilated by $\hat{c}$ and $\hat{d}^+$. The self-consistent equation can be solved explicitly for large U and gives $\Delta = U/2$, which implies that the Mean Field staggered magnetization is 1/2 in that limit.

We define Matsubara correlation functions as

$$\chi^{\mu\nu}(q, q', \tau) = \frac{i}{N} \langle T S^\mu(q, \tau) S^\nu(-q', 0) \rangle \quad (\text{A6})$$

It is straightforward to obtain from them the retarded non-interacting response functions

$$\begin{aligned} \chi_0^{00}(q, q', w) &= \delta_{q,q'} \sum_k' \left(1 - \frac{\Delta^2 + \epsilon_k \epsilon_{k+q}}{E_k E_{k+q}} \right) \left\{ \frac{1}{w + E_{k+q} + E_k + i\delta} - \frac{1}{w - E_{k+q} - E_k + i\delta} \right\} \\ \chi_0^{zz}(q, q', w) &= \frac{\chi_0^{00}(q, q', w)}{4} \\ \chi_0^{+-}(q, q', w) &= \frac{\delta_{q,q'}}{2} \sum_k' \left(1 + \frac{\Delta^2 - \epsilon_k \epsilon_{k+q}}{E_k E_{k+q}} \right) \left\{ \frac{1}{w + E_{k+q} + E_k + i\delta} - \frac{1}{w - E_{k+q} - E_k + i\delta} \right\} \\ \chi_Q^{+-}(q, q', w) &= \frac{\delta_{q,q'+Q} \Delta}{2} \sum_k' \left(\frac{1}{E_k} + \frac{1}{E_{k+q}} \right) \left\{ \frac{1}{w + E_{k+q} + E_k + i\delta} + \frac{1}{w - E_{k+q} - E_k + i\delta} \right\} \quad (\text{A7}) \end{aligned}$$

The RPA-SDW expressions are now given by

$$\begin{aligned} \chi_{RPA}^{00} &= \frac{\chi_0^{00}}{1 + \frac{U}{2} \chi_0^{00}} \\ \chi_{RPA}^{zz} &= \frac{\chi_0^{zz}}{1 - 2U \chi_0^{zz}} \\ \chi_{RPA}^{+-} &= \begin{pmatrix} \chi_o^{+-}(q, w) & \chi_Q^{+-}(q, w) \\ \chi_Q^{+-}(q, w) & \chi_0^{+-}(q + Q, w) \end{pmatrix} \begin{pmatrix} 1 - U \chi_o^{+-}(q, w) & -U \chi_Q^{+-}(q, w) \\ -U \chi_Q^{+-}(q, w) & 1 - U \chi_0^{+-}(q + Q, w) \end{pmatrix}^{-1} \quad (\text{A8}) \end{aligned}$$

The poles of χ_{RPA}^{+-} give the spin-wave excitation spectrum in the RPA-SDW approximation. They are located around the antiferromagnetic, $q_0 = \pi$, and ferromagnetic, $q_0 = 0$, wave-vectors. Both have the same velocity, but different weights. Expanding the denominator for small $q' = q_0 - \pi$ and w , the spin-wave velocity, v_s , in the $q' \rightarrow 0$ limit can be written as

$$\frac{v_s}{t} = 4 \sqrt{\frac{y z U}{x}} \quad (\text{A9})$$

where x , y and z are given by

$$\begin{aligned}
x &= \sum_k' \frac{1}{E_k^3} \\
y &= \sum_k' \frac{3 \sin^2(k) - 1}{E_k^3} - 12 \sum_k' \frac{\cos^2(k) \sin^2(k)}{E_k^5} \\
z &= \sum_k' \frac{\cos^2(k)}{E_k^3}
\end{aligned} \tag{A10}$$

This expression for the spin-wave velocity gives the correct asymptotics for a half-filled band,

$$\begin{aligned}
\frac{v_s}{t} &\rightarrow \sqrt{d} J \quad U \rightarrow \infty \\
&\rightarrow 0 \quad U \rightarrow 0
\end{aligned}$$

where $J = \frac{4t^2}{U}$ is the exchange integral.

The summations over the restricted Brillouin zone can be performed analytically for large U . A straightforward yet tedious calculation gives the following expression for the transverse correlation function

$$\chi_{RPA}^{+-}(q, w) = \frac{1}{U} \frac{w^2 - w_q^2 + 4(1 - \cos(q))}{2w_q} \left\{ \frac{1}{w + w_q + i\delta} - \frac{1}{w - w_q + i\delta} \right\} \tag{A11}$$

where $w_q = v_s |\sin(q)|$ is the spectrum of spin waves. The imaginary part of the transverse response function gives the spectral density of these low-energy excitations:

$$\chi_{RPA}^{+-}{}''(q, w) = \frac{1}{2} \tan\left(\frac{q}{2}\right) \delta(w - w_q) \tag{A12}$$

Notice that $\chi_{RPA}^{+-}{}''(q, w)$ has a zero mode at $q = 0$ with vanishing weight and another at $q = \pi$, whose weight diverges. The real part of the response function at zero energy

$$\chi_{RPA}^{+-}{}''(q, 0) \simeq \frac{2}{U v_s^2} \frac{1}{q^2} \tag{A13}$$

is related to the transverse correlation function C^{+-} .

The RPA-SDW staggered magnetization can be written as

$$\langle S^z \rangle = \langle \hat{m}_i \rangle = \frac{1}{2N} \sum_{k, \sigma} \sigma \int_{-\infty}^{\infty} \frac{dw}{\pi} A_{\sigma}(k, k + Q, w) n_F(w) = \frac{1}{N} \sum_k \int_{-\infty}^0 \frac{dw}{2\pi} A_{\uparrow}(k, k + Q, w) \tag{A14}$$

where the spectral function $A_\sigma(k, k', w)$ is obtained from Dyson's equation. The RPA-SDW self-energy is given by the matrix equation

$$\tilde{\Sigma}_\sigma(k, w) = \frac{U^2}{\beta} \sum_{k', w_B} \tilde{G}_\sigma^0(k + k', w + w_B) \tilde{\chi}_{-\sigma}(k', w_B) \quad (\text{A15})$$

where only the transverse and longitudinal spin plus charge fluctuation channels are taken into account¹². It is important to realize that RPA fluctuations reduce the staggered magnetization from its Mean Field value, $\langle S^z \rangle = \Delta/U$. In particular, for large values of U , Mean Field Theory yields $\langle S^z \rangle = 1/2$ while RPA-SDW gives

$$\langle S^z \rangle = \frac{1}{2} \left\{ 1 - 2 \sum_k \frac{1 - \sin(k)}{|\sin(k)|} \right\} \rightarrow -\infty \quad (\text{A16})$$

This result implies that one can have a charge insulating spin disordered state, or *spin liquid*, within SDW-RPA theory, i.e.: $\langle S^z \rangle \leq 0$ even though $\Delta \neq 0$.

Finally, the expectation value of $\langle S^2 \rangle$ is defined in terms of the dynamic correlation function as

$$\langle S^2 \rangle - \langle S^z \rangle^2 = \sum_q \int_{-\infty}^{\infty} \frac{dw}{2\pi} C(q, w) \quad (\text{A17})$$

which, in turn, is related with the dynamic response function through the fluctuation-dissipation theorem which reads at $T = 0$:

$$\langle S^2 \rangle - \langle S^z \rangle = \sum_q \int_0^\infty \frac{dw}{\pi} (\chi^{zz''}(q, w) + \chi^{+-''}(q, w)) \quad (\text{A18})$$

The double prime denotes again the imaginary part of the function and the summation is now performed over the whole Brillouin zone. It is interesting to notice that the Mean Field approach gives both the weak- and strong-coupling limits correctly

$$\begin{aligned} \langle S^2 \rangle &= S^{zz}(q, w) + S^{+-}(q, w) + \langle S^z \rangle^2 = \frac{1}{8} + \frac{1}{4} = \frac{3}{8} & U = 0 \\ &= 0 + \frac{1}{2} + \frac{1}{4} & U = \infty \end{aligned}$$

APPENDIX B: BETHE ANSATZ RESULTS

The Hubbard model was solved by Lieb and Wu (LW)¹⁵ in 1968. Since then, a number of authors have completed LW's work and obtained the different physical magnitudes^{20,30-33}. This appendix is devoted to display the integral equations, the charge energy gap, $\langle S^2 \rangle$ and v_s as obtained by Shiba³⁰ and Ovchinnikov²⁰.

The integral equations at zero field and half-filling for the distribution of the k-rapidities are given by

$$\begin{aligned}\rho(k) &= \frac{1}{2\pi} + \frac{2}{\pi} \cos(k) \int_{-\pi}^{\pi} dk' \rho(k') S(2(\sin(k) - \sin(k')), u) \\ \partial_u \rho(k) &= \frac{2}{\pi u} \cos(k) \int_{-\pi}^{\pi} dk' \rho(k') R(2(\sin(k) - \sin(k')), u) + \\ &\quad + \frac{2}{\pi} \cos(k) \int_{-\pi}^{\pi} dk' \partial_u \rho(k') \{S(2(\sin(k) - \sin(k')), u) - S(2 \sin(k), u)\}\end{aligned}\tag{B1}$$

where the kernels S and R are

$$\begin{aligned}S(x, u) &= \sum_1^{\infty} (-1)^{n+1} \frac{nu}{x^2 + (nu)^2} \\ R(x, u) &= \sum_1^{\infty} (-1)^{n+1} \frac{(x^2 - (nu)^2)nu}{(x^2 + (nu)^2)^2}\end{aligned}$$

The charge energy gap is given by

$$\frac{\Delta}{t} = 16 u \int_1^{\infty} dx \frac{\sqrt{x^2 - 1}}{\sinh(\frac{2\pi x}{u})}\tag{B2}$$

while the velocity of spin waves in the $q \rightarrow 0$ limit has the following expression

$$\frac{v_s}{t} = \frac{2 I_1(\frac{2\pi}{u})}{I_0(\frac{2\pi}{u})}\tag{B3}$$

where I_n are Bessel functions of imaginary argument. Finally,

$$\langle S^2 \rangle = \frac{3}{4} + 3 \int_{-\pi}^{\pi} dk (1 + \cos(k)) \partial_u \rho(k)\tag{B4}$$

REFERENCES

- ¹ B. Keimer, N. Belk, R. J. Birgueneau, A. Cassanho, C. Y. Chen, M. Greven, M. A. Kastner, A. Aharony, Y. Endoh, R. W. Erwin and G. Shirane, Phys. Rev. B **46**, 14034 (1992)
- ² S. Chakravarty, B. I. Halperin and D. R. Nelson, Phys. Rev. B **39**, 2344 (1989)
- ³ A. V. Chubukov, S. Sachdev and J. Ye, Phys. Rev. B **49**, 11919 (1994)
- ⁴ P. W. Anderson, Science **235**, 1196 (1987)
- ⁵ A. Sokol and D. Pines, Phys. Rev. Lett. **71**, 2813 (1993); A. J. Millis, H. Monien and D. Pines, Phys. Rev. B **42**, 167 (1990)
- ⁶ A. V. Chubukov and K. Musaelian, unpublished
- ⁷ B. I. Shraiman and E. D. Siggia, Phys. Rev. B **42**, 2485 (1990)
- ⁸ A. J. Millis, Phys. Rev. Lett. **71**, 3614 (1993)
- ⁹ A. V. Chubukov and S. Sachdev, Phys. Rev. Lett. **71**, 3615 (1993)
- ¹⁰ H. J. Schulz, Phys. Rev. Lett. **64**, 1445 (1990)
- ¹¹ R. E. Walstedt, B. S. Shastry and S.-W. Cheong, Phys. Rev. Lett. **72**, 3610 (1994)
- ¹² J. R. Schrieffer, X. G. Wen and S. C. Zhang, Phys. Rev. B **39**, 11663 (1989)
- ¹³ A. V. Chubukov and D. M. Frenkel, Phys. Rev. B **46**, 11884 (1992)
- ¹⁴ M. Randeria, N. Trivedi, A. Moreo and R. Scalettar, Phys. Rev. Lett. **69**, 2001 (1992)
- ¹⁵ E. Lieb and F. Y. Wu, Phys. Rev. Lett. **20**, 1445 (1968)
- ¹⁶ H. Shiba, Prog. Theor. Phys. **48**, 2171 (1972); V. J. Emery, Phys. Rev. B **14**, 2989 (1976)
- ¹⁷ D. C. Mattis, The theory of Magnetism I, Springer Series in Solid-State Sciences, Springer-Verlag (1981)

- ¹⁸ N. D. Mermin and H. Wagner, Phys. Rev. Lett. **17**, 1144 (1966)
- ¹⁹ This argument does not hold for some systems, whose low energy excitation spectrum is exhausted by topological solitons.
- ²⁰ A. A. Ovchinnikov, Zh. Eksp. Teor. Fiz. **57**, 2137 (1969) [Soviet Physics JETP, **30**, 1160 (1970)]
- ²¹ F. D. M. Haldane, Phys. Rev. Lett. **61**, 1029 (1988)
- ²² J. Igarashi, Phys. Rev. B **46**, 10763 (1992)
- ²³ J. P. Rodriguez, J. Bonca and J. Ferrer, Phys. Rev B, August 1994; J. Ferrer, Phys. Rev. B **47**, 8769 (1993)
- ²⁴ P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964); A. I. Larkin and Y. N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **47**, 1136 (1964)
- ²⁵ Heavy Fermion superconductors might show this kind of pairing
- ²⁶ P. Nozieres and S. Schmitt-Rink, J. Low Temp. Phys. **59**, 195 (1985)
- ²⁷ C. A. R. Sá de Melo, M. Randeria and J. R. Engelbrecht, Phys. Rev. Lett. **71**, 3202 (1993)
- ²⁸ B. I. Halperin and P. C. Hohenberg, Phys. Rev. **177**, 952 (1969)
- ²⁹ B. D. Josephson, Phys. Lett. **21**, 608 (1966)
- ³⁰ H. Shiba, Phys. Rev. B **6**, 930 (1972); Prog. Theor. Phys. **43**, 1619 (1970)
- ³¹ F. Woyrnarovich, J. Phys. C **16**, 5293 (1983)
- ³² H. Frahm and V. E. Korepin, Phys. Rev. B **42**, 10553 (1990); **43**, 5653 (1991)
- ³³ M. Ogata and H. Shiba, Phys. Rev. B **41**, 2326 (1990)

FIGURES

FIG. 1. Charge gap, 2Δ as a function of U . Solid line is the bethe ansatz exact answer; dashed line is the RPA-SDW Mean Field result

FIG. 2. (a) χ_{RPA}^{+-} as a function of q and w for $U = 20$; (b) $\chi_{RPA}^{zz}(q, w)$ for the same value of U

FIG. 3. Spin-wave velocity, v_s , versus U . Solid line is the exact result; dashed line is RPA's answer. Notice that v_s tends to 0 as U goes to 0, and shows a maximum at $U \sim 1$

FIG. 4. $\langle S^2 \rangle$ versus U . Solid line is the Bethe ansatz result and dashed line is Mean Field's answer

FIG. 5. Characteristic Temperature of the different phenomena as a function of U . Solid line is the Binding Temperature, T_b ; dashed line is the Temperature, T_{corr} , at which correlations begin to build up; dotted line is the transition Temperature to Long Range Order, T_c